



**First Semester B.E. Degree Examination, Dec.2016/Jan.2017**  
**Engineering Mathematics – I**

Time: 3 hrs.

Max. Marks:100

Note: Answer any FIVE full questions, choosing at least two from each part.

**PART – A**

1 a. Choose the correct answers for the following : (04 Marks)

i) If  $y = \frac{1}{2x+1}$  then the 10<sup>th</sup> derivative of y is

A)  $\frac{2^{10}10!}{(-2x+1)^{11}}$       B)  $\frac{2^{10}10!}{(2x+1)^{11}}$       C)  $\frac{2^{10}10!}{(2x-1)^{11}}$       D)  $\frac{2^{10}10!}{(2x+1)^{-11}}$

ii) If  $y = \sin 2x$  then  $y_n$  is

A)  $2^n \sin\left(2x + \frac{n\pi}{2}\right)$       B)  $2^n \cos\left(2x + \frac{n\pi}{2}\right)$       C)  $2^n \sin\left(2x - \frac{n\pi}{2}\right)$       D) none of these

iii) If  $f(x)$  is continuous in  $[a, b]$ , differentiable in  $(a, b)$  and  $f(a) = f(b)$ , then there exist atleast one point  $c \in (a, b)$  such that  $f'(c)$  is equal to

A) 0      B) -1      C)  $\frac{f(b)-f(a)}{b-a}$       D)  $\frac{f(b)-f(a)}{a-b}$

iv) Maclaurin's series expansion of  $e^x$  is

A)  $1 + 2x + \frac{x^2}{2} + \dots$       B)  $1 + x + \frac{x^2}{2} + \dots$       C)  $1 - 2x + \frac{x^2}{2} - \dots$       D)  $1 - x + \frac{x^2}{2} - \dots$

b. If  $\cos^{-1}\left(\frac{y}{b}\right) = \log\left(\frac{x}{n}\right)^p$ , prove that  $x^2 y_{n+2} + (2n+1)xy_{n+1} + (n^2 + p^2)y_n = 0$ . (06 Marks)

c. State Rolle's theorem and verify the theorem for the function  $f(x) = \log\left(\frac{x^2 + ab}{x(a+b)}\right)$  in  $[a, b]$ ,  $b > a > 0$ . (05 Marks)

d. Find the Maclaurin's series expansion of  $\log(1 + e^x)$  upto the term containing  $x^4$ . (05 Marks)

2 a. Choose the correct answers for the following : (04 Marks)

i) The value of  $\lim_{x \rightarrow 0} \frac{a^x - b^x}{x}$  is

A)  $\log\left(\frac{b}{a}\right)$       B)  $\log\left(\frac{a}{b}\right)$       C)  $\log(a-b)$       D) 1

ii) Angle between radius vector and tangent to the curve  $r = a \sin \theta$  is

A)  $\theta$       B)  $-\theta$       C)  $\frac{\pi}{2} - \theta$       D)  $\frac{\pi}{2} + \theta$

iii) The radius of curvature of any point on the curve  $x = a \cos \theta$  and  $y = a \sin \theta$  is

A)  $a \sin \theta$       B)  $\theta$       C)  $\frac{\theta}{2}$       D)  $a$

iv) The derivative of arc length  $\frac{ds}{d\theta}$  for the polar curve  $r = f(\theta)$  is

A)  $\sqrt{r^2 + \frac{d^2r}{d\theta^2}}$       B)  $\sqrt{r + \left(\frac{dr}{d\theta}\right)^2}$       C)  $\sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2}$       D)  $\sqrt{r^2 + \left(\frac{d\theta}{dr}\right)^2}$

b. Evaluate the following:

i)  $\lim_{x \rightarrow 0} (1+x)^{1/x}$       ii)  $\lim_{x \rightarrow 0} \frac{(1+x)^{1/x} - e}{x}$       (06 Marks)

c. For the curve  $y = \frac{ax}{a+x}$ , if  $\rho$  is the radius of curvature at any point  $(x, y)$ , show that

$$\left(\frac{2\rho}{a}\right)^2 = \left(\frac{y}{x}\right)^2 + \left(\frac{x}{y}\right)^2. \quad (05 \text{ Marks})$$

d. Find the angle of intersection of the following pair of curves  $r = a \log \theta$ ,  $r = \frac{a}{\log \theta}$ .      (05 Marks)

3 a. Choose the correct answers for the following :      (04 Marks)

i) If  $u = ax^2 + by^2 + abxy$ , then  $\frac{\partial^3 u}{\partial x^2 \partial y}$  is  
 A) zero      B)  $a + b + ab$       C)  $ab$       D) none of these

ii) If  $u = x^4 y^5$ , where  $x = t^2$  and  $y = t^3$ , then  $\frac{du}{dt}$  is  
 A)  $22 t^{23}$       B)  $20 t^{19}$       C)  $9 t^8$       D)  $23 t^{22}$

iii) If  $x = r \cos \theta$  and  $y = r \sin \theta$  then  $\frac{\partial(x, y)}{\partial(r, \theta)}$  is  
 A)  $r^2 \sin 2\theta$       B)  $r^2$       C)  $r$       D)  $r \sin 2\theta$

iv) The necessary condition for  $u = f(x, y)$  to be extremal is  
 A)  $u_x \neq 0, u_y \neq 0$       B)  $u_x = 0, u_y = 0$       C)  $u_x > 0, u_y > 0$       D)  $u_x < 0, u_y < 0$

b. If  $u = x + 3y^2 - z^3$ ,  $v = 4x^2 yz$ ,  $w = 2z^2 - xy$ , prove that  $\frac{\partial(u, v, w)}{\partial(x, y, z)}$  at  $(1, -1, 0)$  is 20.      (06 Marks)

c. If  $z = \cos(x + ay) + \sin(x - ay)$  prove that  $\frac{\partial^2 z}{\partial y^2} = a^2 \frac{\partial^2 z}{\partial x^2}$ .      (05 Marks)

d. The deflection at the centre of a rod of length  $\ell$  and diameter  $d$ , supported at its ends and located at the centre a weight  $w$ , which varies as  $w\ell^3 d^{-4}$ . Determine the percentage increase in  $w, \ell$  and  $d$  of 5, 4 and 3 respectively.      (05 Marks)

4 a. Choose the correct answers for the following :      (04 Marks)

i) If  $\vec{F} = 3x^2 \hat{i} - xy \hat{j} + (a-3)xz \hat{k}$  is solenoidal, then  $a$  is  
 A) 0      B) -2      C) 2      D) 3

ii) If  $\vec{A} = x^2 \hat{i} + y^2 \hat{j} + z^2 \hat{k}$ , then  $\text{curl } \vec{A}$  is given by  
 A)  $2x \hat{i} + 2y \hat{j} + 2z \hat{k}$       B) 0      C)  $\frac{x \hat{i} + y \hat{j} + z \hat{k}}{2}$       D)  $2x + 2y + 2z$

iii) If  $\phi = xy + yz + zx$ , then  $\text{grad } \phi$  at  $(1, 1, 1)$  is  
 A)  $2 \hat{i} + 2 \hat{j} + 2 \hat{k}$       B) 0      C)  $\hat{i} + \hat{j} + \hat{k}$       D)  $3 \hat{i} + 3 \hat{j} + 3 \hat{k}$

iv) The gradient of a scalar field is a  
 A) vector      B) scalar      C) constant      D) none of these

b. If  $\vec{F} = (x + y + z) \hat{i} + \hat{j} - (x + y) \hat{k}$  then show that  $\vec{F} \cdot \text{curl } \vec{F} = 0$ .      (06 Marks)

c. If  $\vec{F} = \nabla(x^3 + y^3 + z^3 - 3xyz)$  then prove that  $\vec{F}$  is irrotational.      (05 Marks)

d. Derive an expression for  $\text{div } \vec{F}$  in orthogonal curvilinear coordinates.      (05 Marks)

**PART – B**

- 5 a. Choose the correct answers for the following : (04 Marks)
- i) The Leibnitz's rule for differentiation under the integral sign is
- A)  $\phi'(y) = \int_a^b \frac{\partial}{\partial y} [f(x, y)] dx$       B)  $\phi'(y) = \int_a^b \frac{\partial}{\partial x \partial y} [f(x, y)] dx$
- C)  $\phi(y) = \int_a^b \frac{\partial}{\partial x} [f(x, y)] dx$       D) none of these
- ii) The value of  $\int_0^{\pi/2} \sin^6 x dx$  is
- A)  $\frac{5\pi}{8}$       B)  $\frac{5\pi}{64}$       C)  $\frac{5\pi}{32}$       D)  $\frac{5\pi}{16}$
- iii) The value of  $\int_0^{\pi/2} \sin^5 x \cos^5 x dx$  is
- A)  $\frac{1}{90}$       B)  $\frac{1}{60}$       C)  $\frac{1}{30}$       D)  $\frac{1}{70}$
- iv) Surface area of a solid of revolution of the curve  $y = f(x)$ , if rotated about x-axis is
- A)  $\int_{x=a}^b 2\pi y dx$       B)  $\int_{x=a}^b 2\pi x dy$       C)  $\int_{x=a}^b 2\pi y ds$       D)  $\int_{x=a}^b 2\pi x ds$
- b. Using the rule of differentiation under the integral sign, evaluate  $\int_0^{\pi} \frac{\log(1 + \alpha \cos x)}{\cos x} dx$ . (06 Marks)
- c. Obtain the reduction formula for  $\int_0^{\pi/2} \cos^n x dx$ . (05 Marks)
- d. Find the area of the Cardioid  $r = a(1 + \cos \theta)$ . (05 Marks)
- 6 a. Choose the correct answers for the following : (04 Marks)
- i) The solution of  $\frac{dy}{dx} + \frac{y}{x} = 0$  is
- A)  $\frac{y}{x} = c$       B)  $\frac{x}{y} = c$       C)  $x - y = c$       D)  $xy = c$
- ii) The orthogonal trajectory of the family of lines  $y = ax$  is
- A)  $x^2 + y^2 = c^2$       B)  $x^2 - y^2 = c^2$       C)  $xy = c$       D)  $\frac{x}{y} = c$
- iii) The solution of the differential equation  $\frac{dy}{dx} = 1 + \frac{y}{x}$  is
- A)  $y = \log x + c$       B)  $y = x \log x + c$       C)  $y = x(\log x + c)$       D) none of these
- iv) The general solution of the differential equation  $(x - y)dx - (x + y)dy = 0$  is
- A)  $\frac{x^2}{2} - y - \frac{y^2}{2} = c$       B)  $\frac{x^2}{2} - y + \frac{y^2}{2} = c$       C)  $\frac{x^2}{2} - yx + \frac{y^2}{2} = c$       D) none of these
- b. Solve  $\frac{dy}{dx} = \frac{y}{x + \sqrt{xy}}$ . (06 Marks)
- c. Solve  $x \log x \frac{dy}{dx} + y = 2 \log x$ . (05 Marks)
- d. Find the orthogonal trajectories of the family  $x^{2/3} + y^{2/3} = a^{2/3}$ .

