

CBCS SCHEME

USN

--	--	--	--	--	--	--	--	--	--

17MAT11

First Semester B.E. Degree Examination, July/August 2021

Engineering Mathematics – I

Time: 3 hrs.

Max. Marks: 100

Note: Answer any FIVE full questions.

- 1 a. Find the n^{th} derivative of $\cos x \cos 3x \cos 5x$. (06 Marks)
b. If $\tan y = x$, then prove that $(1 + x^2)y_{n+2} + 2(n+1)xy_{n+1} + n(n+1)y_n = 0$ (07 Marks)
c. Derive the angle between radius vector and the tangent. (07 Marks)
- 2 a. If $y = a \cos(\log x) + b \sin(\log x)$ then show that $x^2 y_{n+2} + (2n+1)xy_{n+1} + (n^2 + 1)y_n = 0$. (06 Marks)
b. Find the pedal equation of the polar curve $\frac{2a}{r} = (1 + \cos \theta)$. (07 Marks)
c. Find the radius of curvature for the folium of De-Cartes $x^3 + y^3 = 3axy$ at $(3a/2, 3a/2)$. (07 Marks)
- 3 a. Expand $e^{\sin x}$ using Maclaurin's theorem upto the term containing x^4 . (06 Marks)
b. If $U = \log \sqrt{x^2 + y^2 + z^2}$ show that $(x^2 + y^2 + z^2)[U_{xx} + U_{yy} + U_{zz}] = 1$. (07 Marks)
c. If $x = r \sin \theta \cos \phi$, $y = r \sin \theta \sin \phi$, $z = r \cos \theta$ show that $J\left(\frac{x, y, z}{r, \theta, \phi}\right) = r^2 \sin \theta$. (07 Marks)
- 4 a. Expand $\log(1 + \cos x)$ by Maclaurin's series upto the term containing x^4 . (06 Marks)
b. If $u = f\left(\frac{x}{y}, \frac{y}{z}, \frac{z}{x}\right)$, prove that $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} + z \frac{\partial u}{\partial z} = 0$ (07 Marks)
c. If $u = \frac{yz}{x}$, $v = \frac{zx}{y}$, $w = \frac{xy}{z}$ show that $\frac{\partial(u, v, w)}{\partial(x, y, z)} = 4$ (07 Marks)
- 5 a. A particle moves along the curve $x = t^3 - 4t$, $y = t^2 + 4t$, $z = 8t^2 - 3t^3$ where 't' denotes time. Find the component of its acceleration at $t = 2$ along the tangent. (06 Marks)
b. Find $\text{div } \vec{F}$ and $\text{curl } \vec{F}$ where $\vec{F} = \nabla(x^3 + y^3 + z^3 - 3xyz)$. (07 Marks)
c. Prove that $\text{div}(\text{curl } \vec{A}) = 0$. (07 Marks)
- 6 a. A particle moves along a curve whose parametric equations are $x = e^{-t}$, $y = 2\cos 3t$, $z = 2\sin 3t$. Find the velocity and acceleration at any time 't' and also their magnitudes at $t = 0$. (06 Marks)
b. If $\vec{A} = xz^3\mathbf{i} - 2x^2yz\mathbf{j} + 2yz^4\mathbf{k}$ find $\nabla \cdot \vec{A}$ and $\nabla \cdot (\nabla \times \vec{A})$ at the point $(1, -1, 1)$. (07 Marks)
c. Find the directional derivatives of $\phi = \frac{xz}{x^2 + y^2}$ at $(1, -1, 1)$ along $\mathbf{i} - 2\mathbf{j} + \mathbf{k}$. (07 Marks)

- 7 a. Obtain the reduction formula for $\int \cos^n x$ and hence evaluate $\int_0^{\pi/2} \cos^n x dx$ (06 Marks)
- b. Solve $(x^2 + y^2 + x) dx + xy dy = 0$ (07 Marks)
- c. Find the orthogonal trajectories of the family of curves $\frac{x^2}{a^2} + \frac{y^2}{b^2 + \lambda} = 1$ where λ is a constant. (07 Marks)
- 8 a. Evaluate $\int_0^{\pi/6} \cos^4 3x \sin^2 6x$ using reduction formula. (06 Marks)
- b. Solve $\frac{dy}{dx} + \frac{x+3y-4}{3x+9y-2} = 0$ (07 Marks)
- c. Find the orthogonal trajectories of the family of curves $\left(r + \frac{k^2}{r}\right) \cos \theta = a$, 'a' being parameter. (07 Marks)
- 9 a. Solve the following system of equations by Gauss-Seidel method to obtain the final solution correct to three decimal places.
 $x + y + 54z = 110$, $27x + 6y - z = 85$, $6x + 15y + 2z = 72$. (06 Marks)
- b. Reduce the matrix $A = \begin{bmatrix} -1 & 3 \\ -2 & 4 \end{bmatrix}$ to diagonal form. (07 Marks)
- c. Find the numerically largest eigen value and the corresponding eigen vector of the following matrix using power method
 $A = \begin{bmatrix} 4 & 1 & -1 \\ -2 & 3 & -1 \\ -2 & 1 & 5 \end{bmatrix}$ taking initial approximation as $[1 \ 0.8 \ -0.8]^T$ (07 Marks)
- 10 a. Solve by Gauss elimination method $2x + y + 4z = 12$, $4x + 11y - z = 33$, $8x - 3y + 2z = 20$. (06 Marks)
- b. Diagonalize the matrix $\begin{bmatrix} -19 & 7 \\ -42 & 16 \end{bmatrix}$. (07 Marks)
- c. Using Rayleigh's power method find the numerically largest eigen value and the corresponding eigen vector of the matrix
 $A = \begin{bmatrix} 25 & 1 & 2 \\ 1 & 3 & 0 \\ 2 & 0 & -4 \end{bmatrix}$ by taking $[1 \ 0 \ 0]^T$ as the initial vector. (07 Marks)

CMRIT LIBRARY
BANGALORE - 560 037
