

VISVESVARAYA TECHNOLOGICAL UNIVERSITY
BELAGAVI



MATHEMATICS HANDBOOK

I and II Semester BE Program

Effective from the academic year 2022-2023





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI MATHEMATICS HANDBOOK

Derivatives of some standard functions:

$$\frac{d}{dx}(c) = 0$$

$$\frac{d}{dx}(x^n) = n x^{n-1}$$

$$\frac{d}{dx}(e^x) = e^x$$

$$\frac{d}{dx}(a^x) = a^x \log a$$

$$\frac{d}{dx}(\sin x) = \cos x$$

$$\frac{d}{dx}(\cos x) = -\sin x$$

$$\frac{d}{dx}(\tan x) = \sec^2 x$$

$$\frac{d}{dx}(\cot x) = -\operatorname{cosec}^2 x$$

$$\frac{d}{dx}(\sec x) = \sec x \tan x$$

$$\frac{d}{dx}(\operatorname{cosec} x) = -\operatorname{cosec} x \cot x$$

$$\frac{d}{dx}(\log x) = \frac{1}{x}$$

$$\frac{d}{dx}(\log_a x) = \frac{\log_a e}{x}$$

$$\frac{d}{dx}(\sin^{-1} x) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\cos^{-1} x) = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$$

$$\frac{d}{dx}(\cot^{-1} x) = \frac{-1}{1+x^2}$$

$$\frac{d}{dx}(\sinh x) = \cosh x$$

$$\frac{d}{dx}(\cosh x) = \sinh x$$

$$\frac{d}{dx}(\tanh x) = \sec h^2 x$$

$$\frac{d}{dx}(\coth x) = -\operatorname{cosech}^2 x$$

$$\frac{d}{dx}(\sec hx) = -\sec hx \tanh x$$

$$\frac{d}{dx}(\operatorname{cosech} x) = -\operatorname{cosech} x \coth x$$

Rules of Differentiation:

$$\frac{d}{dx}(cu) = c \frac{du}{dx}$$

$$\frac{d}{dx}(uv) = u \frac{d}{dx}(v) + v \frac{d}{dx}(u)$$

$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \frac{d}{dx}(u) - u \frac{d}{dx}(v)}{v^2}$$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

Parametric differentiation:

If $x = x(t)$ & $y = y(t)$ then $\frac{dy}{dx} = \frac{\left(\frac{dy}{dt}\right)}{\left(\frac{dx}{dt}\right)}$

Chain Rule:

If $y = f(u)$ & $u = g(x)$ then $\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx}$

Integrals of some standard functions:

(Constant of Integration C to be added in all the integrals)

$$\int x^n dx = \frac{x^{n+1}}{n+1} \quad \int \frac{1}{x} dx = \log x$$

$$\int \log x dx = x \log x - x, x \neq 0 \quad \int k dx = k x$$

$$\int e^x dx = e^x \quad \int a^x dx = \frac{a^x}{\log a}$$

$$\int \sin x dx = -\cos x \quad \int \cos x dx = \sin x$$

$$\int \tan x dx = \log(\sec x) \quad \int \cot x dx = \log(\sin x)$$

$$\int \sec x dx = \log(\sec x + \tan x) \quad \int \cos ec x dx = \log(\cos ec x - \cot x)$$

$$\int \sec^2 x dx = \tan x \quad \int \cos ec^2 x dx = -\cot x$$

$$\int \sec x \tan x dx = \sec x \quad \int \cos ec x \cot x dx = -\cos ec x$$

$$\int \sinh x dx = \cosh x \quad \int \cosh x dx = \sinh x$$

$$\int \tanh x dx = \log(\cosh x) \quad \int \coth x dx = \log(\sinh x)$$

$$\int \sec h^2 x dx = \tanh x \quad \int \cos ech^2 x dx = -\coth x$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1}(x/a) \quad \int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1}(x/a)$$

$$\int \frac{f'(x)}{f(x)} dx = \log(f(x)) \quad \int \frac{f'(x)}{\sqrt{f(x)}} dx = 2\sqrt{f(x)}$$

$$\int e^{ax} \cos bx dx = \frac{e^{ax}}{a^2 + b^2} [a \cos bx + b \sin bx]$$

$$\int e^{ax} \sin bx dx = \frac{e^{ax}}{a^2 + b^2} [a \sin bx - b \cos bx]$$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

$$\int_0^a f(x) dx = \int_0^a f(a-x) dx$$

$$\int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$\int_{-a}^a f(x) dx = \begin{cases} 2 \int_0^a f(x) dx & \text{if } f(x) \text{ is even function} \\ 0 & \text{if } f(x) \text{ is odd function} \end{cases}$$

$$\int_0^{2a} f(x) dx = \begin{cases} 2 \int_0^a f(x) dx, & \text{if } f(2a-x) = f(x) \\ 0, & \text{if } f(2a-x) = -f(x) \end{cases}$$

Integration by parts:

$$\int u(x)v(x) dx = u(x) \left(\int v(x) dx \right) - \int \frac{d}{dx}(u(x)) \left(\int v(x) dx \right) dx$$

Bernoulli's rule of integration:

If the 1st function is a polynomial and integration of 2nd function is known. Then

$$\int u(x)v(x) dx = u \int v dx - u' \int \int v dx dx + u'' \int \int \int v dx dx dx - u''' \int \int \int \int v dx dx dx dx + \dots \dots \dots$$

Where dashes denote the differentiation of u .

Or

$$\int u(x)v(x) dx = u \cdot v_1 - u' \cdot v_2 + u'' \cdot v_3 - u''' \cdot v_4 + \dots \dots \dots$$

Where dashes denote the differentiation of u , v_k denotes the integration of v , k times with respect to x .

Vector calculus formulae:

Position vector $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$

Magnitude $|\vec{r}| = \sqrt{x^2 + y^2 + z^2}$

Dot product of unit vectors $\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1$ and $\hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0$

Cross product of unit vectors $\hat{i} \times \hat{i} = \hat{j} \times \hat{j} = \hat{k} \times \hat{k} = 0$ and $\hat{i} \times \hat{j} = \hat{k}, \hat{j} \times \hat{k} = \hat{i}, \hat{i} \times \hat{k} = -\hat{j}$

Angle between two vectors $\cos \theta = \frac{\vec{A} \cdot \vec{B}}{|\vec{A}| |\vec{B}|}$

Unit vector $\hat{A} = \frac{\vec{A}}{|\vec{A}|}$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

Velocity $\vec{V} = \frac{d\vec{s}}{dt}$

Acceleration $\vec{a} = \frac{d^2\vec{s}}{dt^2}$

For any vectors $\vec{A} = (a_1 i + b_1 j + c_1 k)$, $\vec{B} = (a_2 i + b_2 j + c_2 k)$ & $\vec{C} = (a_3 i + b_3 j + c_3 k)$

Dot product of two vectors $\vec{A} \cdot \vec{B} = a_1 a_2 + b_1 b_2 + c_1 c_2$

Cross product of two vectors $\vec{A} \times \vec{B} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$

Scalar triple product $\vec{A} \cdot (\vec{B} \times \vec{C}) = (\vec{A} \times \vec{B}) \cdot \vec{C} = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$

Trigonometric formulae:

- **Identities**

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$1 + \tan^2 \theta = \sec^2 \theta$$

$$1 + \cot^2 \theta = \operatorname{cosec}^2 \theta$$

- **Compound angle formulae**

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A-B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A+B) = \cos A \cos B - \sin A \sin B$$

$$\cos(A-B) = \cos A \cos B + \sin A \sin B$$

$$\tan(A+B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\tan(A-B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

- **Transformation formulae**

$$\sin A \cos B = \frac{1}{2} [\sin(A+B) + \sin(A-B)],$$



$$\cos A \sin B = \frac{1}{2} [\sin(A+B) - \sin(A-B)]$$

$$\cos A \cos B = \frac{1}{2} [\cos(A+B) + \cos(A-B)],$$

$$\sin A \sin B = \frac{1}{2} [\cos(A-B) - \cos(A+B)]$$

$$\sin C + \sin D = 2 \sin\left(\frac{C+D}{2}\right) \cos\left(\frac{C-D}{2}\right),$$

$$\sin C - \sin D = 2 \sin\left(\frac{C-D}{2}\right) \cos\left(\frac{C+D}{2}\right)$$

$$\cos C + \cos D = 2 \cos\left(\frac{C+D}{2}\right) \cos\left(\frac{C-D}{2}\right),$$

$$\cos C - \cos D = -2 \sin\left(\frac{C+D}{2}\right) \sin\left(\frac{C-D}{2}\right)$$



VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

- **Multiple angle formulae**

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = \cos^2 A - \sin^2 A$$

$$\sin^2 A = \frac{(1 - \cos 2\theta)}{2}$$

$$\sin^3 A = \frac{1}{4} [3 \sin A - \sin 3A]$$

$$\sin A = 2 \sin(A/2) \cos(A/2)$$

$$\cos A = \cos^2(A/2) - \sin^2(A/2)$$

$$\cos^2 A = \frac{(1 + \cos 2\theta)}{2}$$

$$\cos^3 A = \frac{1}{4} [3 \cos A + \cos 3A]$$

Hyperbolic and Euler's formulae

$$\sinh x = \frac{e^x - e^{-x}}{2}$$

$$\cosh^2 \theta - \sinh^2 \theta = 1$$

$$\cosh x = \frac{e^x + e^{-x}}{2}$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$\sin x = \frac{e^{ix} - e^{-ix}}{2i}$$

$$\cos x = \frac{e^{ix} + e^{-ix}}{2}$$

Logarithmic formulae:

$$\log_e(A B) = \log_e(A) + \log_e(B)$$

$$\log_e\left(\frac{A}{B}\right) = \log_e(A) - \log_e(B)$$

$$\log_e x^n = n \log_e x$$

$$\log_a B = \frac{\log_e B}{\log_e a}$$

$$\log_a a = 1$$

$$\log_a 1 = 0$$

$$\log_e 0 = -\infty$$

Solid geometry formulae:

Distance between two points (x_1, y_1, z_1) and (x_2, y_2, z_2) is $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$

Direction cosines $l = \cos \alpha, m = \cos \beta, n = \cos \gamma$

Direction ratios $l = \frac{a}{\sqrt{a^2 + b^2 + c^2}}, m = \frac{b}{\sqrt{a^2 + b^2 + c^2}}, n = \frac{c}{\sqrt{a^2 + b^2 + c^2}}$.

Direction ratios of a line joining two points $(a, b, c) = (x_2 - x_1, y_2 - y_1, z_2 - z_1)$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

nth Derivatives of standard functions

$$D^n \left[(ax+b)^m \right] = m(m-1)(m-2) \dots (m-n+1)(ax+b)^{m-n} \cdot a^n$$

$$D^n \left[(ax+b)^n \right] = n! a^n$$

$$D^n \left(x^n \right) = n!$$

$$D^n \left(\frac{1}{ax+b} \right) = \frac{(-1)^n n! a^n}{(ax+b)^{n+1}}$$

$$D^n \left[\log(ax+b) \right] = \frac{(-1)^{n-1} (n-1)! a^n}{(ax+b)^n}.$$

$$D^n \left[a^{mx} \right] = a^{mx} (m \log a)^n$$

$$D^n \left(e^{ax} \right) = a^n e^{ax}$$

$$D^n \left[\sin(ax+b) \right] = a^n \sin(ax+b + n\pi/2)$$

$$D^n \left[\cos(ax+b) \right] = a^n \cos(ax+b + n\pi/2)$$

$$D^n \left[e^{ax} \sin(bx+c) \right] = (a^2 + b^2)^{n/2} e^{ax} \sin(bx+c + n \tan^{-1}(b/a))$$

$$D^n \left[e^{ax} \cos(bx+c) \right] = (a^2 + b^2)^{n/2} e^{ax} \cos(bx+c + n \tan^{-1}(b/a))$$

Polar coordinates and polar curves:

Angle between radius vector and tangent

$$\tan \phi = r \frac{d\theta}{dr} \quad \text{or} \quad \cot \phi = \frac{1}{r} \frac{dr}{d\theta}$$

Angle of intersection of the curves

$$|\phi_1 - \phi_2| = \tan^{-1} \left\{ \left| \frac{\tan \phi_1 - \tan \phi_2}{1 + \tan \phi_1 \tan \phi_2} \right| \right\}$$

Orthogonal condition $|\phi_1 - \phi_2| = \frac{\pi}{2}$ or $\tan \phi_1 \cdot \tan \phi_2 = -1$,

Pedal equation or p-r equation

$$P = r \sin \phi$$

$$\frac{1}{p^2} = \frac{1}{r^2} (1 + \cot^2 \phi) = \frac{1}{r^2} \left(1 + \frac{1}{r^2} \left(\frac{dr}{d\theta} \right)^2 \right)$$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

Derivative of arc length:

$$\text{In Cartesian: } \frac{ds}{dx} = \sqrt{1 + \left(\frac{dy}{dx}\right)^2} \quad \& \quad \frac{ds}{dy} = \sqrt{1 + \left(\frac{dx}{dy}\right)^2}$$

$$\text{In Polar: } \frac{ds}{dr} = \sqrt{1 + \frac{1}{r^2} \left(\frac{d\theta}{dr}\right)^2} \quad \& \quad \frac{ds}{d\theta} = \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2}$$

$$\text{In Parametric: } \frac{ds}{dt} = \sqrt{\left(\frac{dx}{dt}\right)^2 + \left(\frac{dy}{dt}\right)^2}$$

$$\sin\theta = r \frac{d\theta}{ds} \quad \& \quad \cos\theta = r \frac{dr}{ds}$$

Radius of curvature

$$\text{In Cartesian form: } \rho = \frac{(1+y_1^2)^{\frac{3}{2}}}{y_2}$$

$$\text{In parametric form: } \rho = \frac{(\dot{x}^2 + \dot{y}^2)^{\frac{3}{2}}}{\dot{x}\dot{y} - \dot{y}\dot{x}}$$

$$\text{In polar form: } \rho = \frac{(r^2 + r_1^2)^{\frac{3}{2}}}{r^2 - rr_1 + 2r_1^2}$$

$$\text{Pedal Equation: } \rho = r \frac{dr}{dp}$$

Indeterminate Forms - L'Hospital's rule:

$$\text{If } f(a) = g(a) = 0, \text{ then } \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$\text{If } f(a) = g(a) = \infty, \text{ then } \lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

$$\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n = e \quad , \quad \lim_{n \rightarrow 0} (1+n)^{\frac{1}{n}} = e$$

$$\lim_{\theta \rightarrow 0} \frac{\sin\theta}{\theta} = 1 \quad , \quad \lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1}$$

Series Expansion:

Taylor's series expansion about the point $x = a$.

$$y(x) = y(a) + \frac{(x-a)}{1!} y'(a) + \frac{(x-a)^2}{2!} y''(a) + \frac{(x-a)^3}{3!} y'''(a) + ..$$

Maclaurin's Series at the point $x = 0$

$$y(x) = y(0) + \frac{x}{1!} y'(0) + \frac{x^2}{2!} y''(0) + \frac{x^3}{3!} y'''(0) + \frac{x^4}{4!} y''''(0) +$$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

Euler's theorem on homogeneous function and Corollary:

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = n u \quad (\text{Theorem})$$

$$x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = n \frac{f(u)}{f'(u)} \quad (\text{Corollary 1})$$

$$x^2 \frac{\partial^2 u}{\partial x^2} + 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} = n(n-1)u \quad (\text{Corollary 2})$$

Composite function:

$$\text{If } z = f(x, y) \text{ and } x = \phi(t), y = \psi(t) \text{ then } \frac{dz}{dt} = \frac{\partial z}{\partial x} \frac{dx}{dt} + \frac{\partial z}{\partial y} \frac{dy}{dt}$$

$$\text{If } z = f(x, y) \text{ and } x = \phi(u, v), y = \psi(u, v)$$

$$\frac{\partial z}{\partial u} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial u} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial u} \quad \& \quad \frac{\partial z}{\partial v} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial v} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial v}$$

$$\text{If } u = f(r, s, t) \text{ and } r = \phi(x, y, z), s = \psi(x, y, z), t = \xi(x, y, z)$$

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial u}{\partial r} \frac{\partial r}{\partial x} + \frac{\partial u}{\partial s} \frac{\partial s}{\partial x} + \frac{\partial u}{\partial t} \frac{\partial t}{\partial x} \\ \frac{\partial u}{\partial y} &= \frac{\partial u}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial u}{\partial s} \frac{\partial s}{\partial y} + \frac{\partial u}{\partial t} \frac{\partial t}{\partial y} \end{aligned}$$

$$\frac{\partial u}{\partial z} = \frac{\partial u}{\partial r} \frac{\partial r}{\partial z} + \frac{\partial u}{\partial s} \frac{\partial s}{\partial z} + \frac{\partial u}{\partial t} \frac{\partial t}{\partial z}$$

Jacobians:

$$J = \frac{\partial(u, v)}{\partial(x, y)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} \end{vmatrix} \quad \text{and} \quad \frac{\partial(u, v, w)}{\partial(x, y, z)} = \begin{vmatrix} \frac{\partial u}{\partial x} & \frac{\partial u}{\partial y} & \frac{\partial u}{\partial z} \\ \frac{\partial v}{\partial x} & \frac{\partial v}{\partial y} & \frac{\partial v}{\partial z} \\ \frac{\partial w}{\partial x} & \frac{\partial w}{\partial y} & \frac{\partial w}{\partial z} \end{vmatrix}$$

Multiple Integrals:

$$\text{Area } A = \iint_A dx dy \text{ - Cartesian form}$$



$$\text{Area } A = \iint_A r dr d\theta \text{ - Polar form}$$



VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI MATHEMATICS HANDBOOK

Volume $V = \iiint_V dx dy dz$ - Cartesian form

Volume $V = \iint_A z dx dy$ - by double integral

$$\text{Gamma function: } \Gamma(n) = \int_0^\infty e^{-x} x^{n-1} dx = 2 \int_0^\infty e^{-t^2} t^{2n-1} dt$$

$$\text{Beta function: } \beta(m, n) = \int_0^1 x^{n-1} (1-x)^{m-1} dx = 2 \int_0^{\frac{\pi}{2}} \sin^{2n-1} \theta \cos^{2m-1} \theta d\theta$$

$$\text{Beta and Gamma relation: } \beta(m, n) = \frac{\Gamma(m)\Gamma(n)}{\Gamma(m+n)} \text{ and } \Gamma(\frac{1}{2}) = \sqrt{\pi}$$

Vector Calculus:

Velocity

$$\vec{v}(t) = \frac{d \vec{r}}{dt}$$

Acceleration

$$\vec{a}(t) = \frac{d \vec{v}}{dt} = \frac{d^2 \vec{r}}{dt^2}.$$

The unit tangent vector

$$\hat{T} = \frac{d \vec{r}}{dt} \Bigg/ \left| \frac{d \vec{r}}{dt} \right|$$

Angle between the tangents

$$\cos \theta = \frac{\vec{T}_1 \cdot \vec{T}_2}{\left| \vec{T}_1 \right| \left| \vec{T}_2 \right|}$$

Component of velocity

$$C.V = \vec{v} \cdot \hat{n} \quad , \text{ Where } \hat{n} \text{ is the unit vector}$$

Component of accelerations

$$C.A = \vec{a} \cdot \hat{n}$$

Tangential component of acceleration

$$T.C.A = \vec{a} \cdot \vec{v} \Bigg/ \left| \vec{v} \right|$$

Normal component of acceleration

$$N.C.A = \left| \vec{a} - (\text{tangential component}) \times \left(\vec{v} \Big/ \left| \vec{v} \right| \right) \right|$$

Gradient of ϕ

$$\text{grad } \phi = \nabla \phi = \frac{\partial \phi}{\partial x} \hat{i} + \frac{\partial \phi}{\partial y} \hat{j} + \frac{\partial \phi}{\partial z} \hat{k}$$

Unit vector normal to the surface

$$\hat{n} = \frac{\nabla \phi}{|\nabla \phi|}$$

Directional Derivative: $D. D = \nabla \phi \cdot \hat{n}$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI MATHEMATICS HANDBOOK

Angle between the surfaces

$$\cos \theta = \frac{\nabla \phi_1 \cdot \nabla \phi_2}{|\nabla \phi_1| |\nabla \phi_2|}$$

Divergence of vector field $\vec{F} = f_1 i + f_2 j + f_3 k$

$$\nabla \cdot \vec{F} = \operatorname{div} \vec{F} = \frac{\partial f_1}{\partial x} + \frac{\partial f_2}{\partial y} + \frac{\partial f_3}{\partial z}$$

Curl of vector field \vec{F}

$$\nabla \times \vec{F} = \operatorname{curl} \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ f_1 & f_2 & f_3 \end{vmatrix}$$

Solenoidal vector field

$$\operatorname{div} \vec{F} = \nabla \cdot \vec{F} = 0$$

Irrational vector field

$$\operatorname{curl} \vec{F} = \nabla \times \vec{F} = 0.$$

List of vector identities

$$\operatorname{curl}(\operatorname{grad} \phi) = \nabla \times \nabla \phi = 0.$$

$$\operatorname{div}(\operatorname{curl} \vec{F}) = \nabla \cdot (\nabla \times \vec{F}) = 0.$$

$$\operatorname{div}(\phi \vec{F}) = \phi \left(\operatorname{div} \vec{F} \right) + \operatorname{grad} \phi \cdot \vec{F}$$

$$\operatorname{curl}(\phi \vec{F}) = \phi \left(\operatorname{curl} \vec{F} \right) + \operatorname{grad} \phi \times \vec{F}$$

Reduction formulae

$$\int_0^{\pi/2} \cos^n x dx = \int_0^{\pi/2} \sin^n x dx = \begin{cases} \frac{(n-1)(n-3)(n-5)\dots\dots 1}{n(n-2)(n-4)\dots\dots 2} \frac{\pi}{2} & \text{when } n \text{ is even} \\ \frac{(n-1)(n-3)(n-5)\dots\dots 3}{n(n-2)(n-4)\dots\dots 1} & \text{when } n \text{ is odd} \end{cases}$$

$$\int_0^{\pi/2} \sin^m x \cos^n x dx = \begin{cases} \frac{(m-1)(m-3)\dots\dots (n-1)(n-3)\dots\dots}{(m+n)(m+n-2)(m+n-4)\dots\dots 2} \frac{\pi}{2} & \text{when } m \text{ and } n \text{ is even} \\ \frac{(m-1)(m-3)\dots\dots (n-1)(n-3)\dots\dots}{(m+n)(m+n-2)(m+n-4)\dots\dots} & \text{other cases} \end{cases}$$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

Differential Equations:

Differential Equations	Solution/ substitution
Linear in y $\frac{dy}{dx} + Py = Q$	$y(I.F.) = \int Q(I.F.)dx + C$
Linear in x $\frac{dx}{dy} + Px = Q$	$x(I.F.) = \int Q(I.F.)dy + C$
Bernoulli's $\frac{dy}{dx} + P y = Q y^n$	divide by y^n and Put $y^{1-n} = z$
Exact differential equation $Mdx + Ndy = 0$ with $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$.	$\int_M dx + \int_{y \text{ constant}} (\text{terms of } N \text{ not containing } x) dy = C$
Not exact differential equation Case 1 : If $Mdx + Ndy = 0$ and $Mdx + Ndy = 0$	$I.F. = \frac{1}{Mx + Ny}, Mx + Ny \neq 0$
Case 2 : If the differential equation is of the form $yg_1(xy)dx + xg_2(xy)dy = 0$	$I.F. = \frac{1}{Mx - Ny} \text{ with } Mx - Ny \neq 0$
Case 3: If $\frac{1}{N} \left(\frac{\partial M}{\partial y} - \frac{\partial N}{\partial x} \right) = f(x) \text{ or } C$	$I.F. = e^{\int f(x)dx} \text{ or } e^{\int Cdx}$
Case 4 : If $\frac{1}{M} \left(\frac{\partial N}{\partial x} - \frac{\partial M}{\partial y} \right) = f(y) \text{ or } C$	$I.F. = e^{\int f(y)dy} \text{ or } e^{\int Cd y}$
Case 5 : If the differential equation is of the form $x^n y^n (mydx + nxdy) + x^p y^q (m' ydx + n' xdy) = 0$	$I.F. = x^h y^k \text{ With}$ $\frac{p+h+1}{m} = \frac{q+k+1}{n} \text{ and } \frac{p'+h+1}{m'} = \frac{q'+k+1}{n'}$
Newton's law of cooling	$T = T_o + C_1 e^{-kt}$

Linear Algebra:

Inverse of a square matrix A

$$A^{-1} = \frac{(adj A)}{|A|}.$$

Rank of a Matrix A

The number of non-zero rows in the echelon form of A is equal to rank of A

Normal Form of a Matrix.

$$\begin{bmatrix} I_r & 0 \\ 0 & 0 \end{bmatrix}$$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI MATHEMATICS HANDBOOK

Gauss-Elimination Method

The system is reduced to upper triangular system from which the unknowns are found by back substitutions.

Gauss-Jordan Method

The system is reduced to diagonal system from which the unknowns are found by back substitutions.

Eigenvalues

Roots of the characteristic equation $|A - \lambda I| = 0$

Eigen Vectors

Non-zero solution $x = x_i$ of $|A - \lambda I| x = 0$

Diagonal form

$$D = P^{-1}AP = \begin{bmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{bmatrix}$$

Computation of power of a square matrix A

$$A^n = P D^n P^{-1}$$

Canonical Form

$$V = \lambda_1 y_1^2 + \lambda_2 y_2^2 + \lambda_3 y_3^2 + \dots + \lambda_n y_n^2$$

Nature, Rank and Index of Quadratic forms:

- **positive-definite** if all the eigenvalues of A are positive.
- **positive-semi definite** if all the eigenvalues of A are non-negative and at least one of the eigenvalues is zero.
- **negative-definite** if all the eigenvalues of A are negative.
- **negative-semi definite** if all the eigenvalues of A are negative and at least one of the eigenvalues is zero.
- **indefinite** if the matrix A has both positive and negative eigenvalues.
- **Rank** the number of non-zero terms
- **Index** the number of positive terms
- **Signature** the number of positive terms minus the number of negative terms





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

Laplace Transforms:

Laplace Transform of Standard Functions

$L\{1\} = \frac{1}{s}$	$L\{t^n\} = \frac{n!}{s^{n+1}}, n=1,2,3.....$ Where n is a positive integer
$L\{e^{at}\} = \frac{1}{s-a}$	$L\{e^{-at}\} = \frac{1}{s+a}$
$L\{\sin at\} = \frac{a}{s^2 + a^2}$	$L\{\cos at\} = \frac{s}{s^2 + a^2}$
$L\{\sinh at\} = \frac{a}{s^2 - a^2}$	$L\{\cosh at\} = \frac{s}{s^2 - a^2}$
$L\{u(t)\} = \frac{1}{s}$	$L\{u(t-a)\} = \frac{e^{-as}}{s}$

Properties of the Laplace transform:

$f(t)$	$L\{f(t)\} = F(s)$
Translation (first Shifting Theorem)	$L\{e^{at} f(t)\} = F(s-a)$
Multiplication by t	$L\{t^n f(t)\} = (-1)^n \frac{d^n}{ds^n} \{F(s)\}$
Time scale	$L\{f(at)\} = \frac{1}{a} F\left(\frac{s}{a}\right)$
Integration	$L\left\{\int_0^t f(\tau) d\tau\right\} = \frac{1}{s} L\{f(t)\}$
Division by t	$L\left\{\frac{f(t)}{t}\right\} = \int_s^\infty F(s) ds.$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

Transform of a Periodic Function: If $f(t)$ is periodic with a period T , then

$$L\{f(t)\} = \frac{1}{(1-e^{-sT})} \int_0^T e^{-st} f(t) dt .$$

Second Shifting Theorem:

If $F(s) = L\{f(t)\}$ and $a > 0$, then $L\{f(t-a)u(t-a)\} = e^{-as} L\{f(t)\}$.

Transforms of the derivatives:

$$L\{f'(t)\} = sL\{f(t)\} - f(0)$$

$$L\{f''(t)\} = s^2 L\{f(t)\} - s f(0) - f'(0)$$

$$L\{f^n(t)\} = s^n L\{f(t)\} - s^{n-1} f(0) - s^{n-2} f'(0) - s^{n-3} f''(0) - \dots - f^{n-1}(0)$$

Inverse Laplace Transform:

Transform	Inverse Transform
$F(s) = L\{f(t)\}$	$f(t) = L^{-1}\{F(s)\}$
$L\{t^n\} = \frac{n!}{s^{n+1}}$ Where n is a positive integer	$\frac{t^{n-1}}{(n-1)!} = L^{-1}\left\{\frac{1}{s^n}\right\}$ Where n is a positive integer
$L\{e^{at}\} = \frac{1}{s-a}$	$e^{at} = L^{-1}\left\{\frac{1}{s-a}\right\}$
$L\{\sinh at\} = \frac{a}{s^2 - a^2}$	$\sinh at = L^{-1}\left\{\frac{a}{s^2 - a^2}\right\}$
$L\{\cosh at\} = \frac{s}{s^2 - a^2}$	$\cosh at = L^{-1}\left\{\frac{s}{s^2 - a^2}\right\}$
$L\{\sin at\} = \frac{a}{s^2 + a^2}$	$\sin at = L^{-1}\left\{\frac{a}{s^2 + a^2}\right\}$
$L\{\cos at\} = \frac{s}{s^2 + a^2}$	$\cos at = L^{-1}\left\{\frac{s}{s^2 + a^2}\right\}$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

Formulae on Shifting rule:

$$\begin{aligned} L[e^{at} \cos bt] &= \frac{s-a}{(s-a)^2+b^2} \\ L[e^{at} \sin bt] &= \frac{b}{(s-a)^2+b^2} \\ L[e^{-at} \cos bt] &= \frac{s+a}{(s+a)^2+b^2} \\ L[e^{-at} \sin bt] &= \frac{b}{(s+a)^2+b^2} \\ L[e^{at} \sinh bt] &= \frac{b}{(s-a)^2-b^2} \\ L[e^{at} \cosh bt] &= \frac{s-a}{(s-a)^2-b^2} \\ L[e^{-at} \cosh bt] &= \frac{s+a}{(s+a)^2-b^2} \\ L[e^{-at} \sinh bt] &= \frac{b}{(s+a)^2-b^2} \end{aligned}$$

$$\begin{aligned} L^{-1}\left[\frac{s-a}{(s-a)^2+b^2}\right] &= e^{at} \cos bt \\ L^{-1}\left[\frac{b}{(s-a)^2+b^2}\right] &= e^{at} \sin bt \\ L^{-1}\left[\frac{s+a}{(s+a)^2+b^2}\right] &= e^{-at} \cos bt \\ L^{-1}\left[\frac{b}{(s+a)^2+b^2}\right] &= e^{-at} \sin b \\ L^{-1}\left[\frac{b}{(s-a)^2-b^2}\right] &= e^{at} \sinh bt \\ L^{-1}\left[\frac{s-a}{(s-a)^2-b^2}\right] &= e^{at} \cosh bt \\ L^{-1}\left[\frac{s+a}{(s+a)^2-b^2}\right] &= e^{-at} \cosh bt \\ L^{-1}\left[\frac{b}{(s+a)^2-b^2}\right] &= e^{-at} \sinh bt \end{aligned}$$

Convolution Theorem:

Let $f(t)$ and $g(t)$ be piecewise continuous on $[0, \infty)$ and $F(s) = L\{f(t)\}$ & $G(s) = L\{g(t)\}$.

Then, $L^{-1}\{F(s)G(s)\} = \int_0^t f(u)g(t-u)du.$

Numerical Methods:

Regula Falsi formula: $x_{k+2} = x_k - \frac{x_{k+1}-x_k}{f(x_{k+1})-f(x_k)}$ for $k = 0, 1, 2, \dots$

Newton-Raphson formula:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)} \quad \text{for } n = 1, 2, 3, \dots$$

Newton's Forward Interpolation formula:

$$y_p = y_0 + p\Delta y_0 + \frac{p(p-1)}{2!} \Delta^2 y_0 + \frac{p(p-1)(p-2)}{3!} \Delta^3 y_0 + \dots$$

Where $p = \frac{x-x_0}{h}$

Newton's Backward Interpolation formula:

$$y_p = y_0 + p\nabla y_0 + \frac{p(p+1)}{2!} \nabla^2 y_0 + \frac{p(p+1)(p+2)}{3!} \nabla^3 y_0 + \dots$$

Where $p = \frac{x-x_n}{h}$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

Newton's General Interpolation formula (Divided difference formula):

$$y = f(x) = y_0 + (x - x_0)[x_0, x_1] + (x - x_0)(x - x_1)[x_0, x_1, x_2] \\ + (x - x_0)(x - x_1)(x - x_2)[x_0, x_1, x_2, x_3] + \dots \dots \dots \\ + (x - x_0)(x - x_1) \dots \dots (x - x_n)[x_0, x_1, \dots, x_n]$$

Lagrange's Interpolation formula:

$$f(x) = \frac{(x - x_1)(x - x_2)(x - x_3) \dots \dots (x - x_n)}{(x_0 - x_1)(x_0 - x_2)(x_0 - x_3) \dots \dots (x_0 - x_n)} y_0 \\ + \frac{(x - x_0)(x - x_2)(x - x_3) \dots \dots (x - x_n)}{(x_1 - x_0)(x_1 - x_2)(x_1 - x_3) \dots \dots (x_1 - x_n)} y_1 \\ + \frac{(x - x_0)(x - x_1)(x - x_3) \dots \dots (x - x_n)}{(x_2 - x_0)(x_2 - x_1)(x_2 - x_3) \dots \dots (x_2 - x_n)} y_2 \\ + \dots \dots \dots \\ + \frac{(x - x_0)(x - x_1)(x - x_2) \dots \dots (x - x_n)}{(x_n - x_0)(x_n - x_1)(x_n - x_2) \dots \dots (x_n - x_{n-1})} y_n$$

Numerical Integration:

Trapezoidal Rule:

$$\int_{x_0}^{x_0+nh} f(x) dx = \frac{h}{2} [(y_0 + y_n) + 2(y_1 + y_2 + \dots + y_{n-1})]$$

Simpson's (1/3)rd rule:

$$\int_{x_0}^{x_0+nh} f(x) dx = \frac{h}{3} [(y_0 + y_n) + 4(y_1 + y_3 + \dots + y_{n-1}) + 2(y_2 + y_4 + \dots + y_{n-2})]$$

Simpson's (3/8)th rule:

$$\int_{x_0}^{x_0+nh} f(x) dx \\ = \frac{3h}{8} [(y_0 + y_n) + 3(y_1 + y_2 + y_4 + y_5 + \dots + y_{n-1}) \\ + 2(y_3 + y_6 + \dots + y_{n-3})]$$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI MATHEMATICS HANDBOOK

Numerical methods for ODE's:

Taylor's series expansion about the point $x = x_0$.

$$y(x) = y(x_0) + \frac{(x-x_0)}{1!} y'(x_0) + \frac{(x-x_0)^2}{2!} y''(x_0) + \frac{(x-x_0)^3}{3!} y'''(x_0) + \dots$$

Taylor's series expansion about the point $x = 0$

$$y(x) = y(0) + \frac{x}{1!} y'(0) + \frac{x^2}{2!} y''(0) + \frac{x^3}{3!} y'''(0) + \frac{x^4}{4!} y''''(0) + \dots \quad \text{Euler's Method:}$$

$$y_{n+1}^E = y_n + hf(x_n, y_n), \quad n = 0, 1, 2, 3, \dots$$

Modified Euler's Method:

$$y_{n+1} = y_n + \frac{h}{2} [f(x_n, y_n) + f(x_{n+1}, y_{n+1}^E)], \quad n = 0, 1, 2, 3, \dots$$

Runge-Kutta Method

Step1: Find $k_1 = hf(x_n, y_n); \quad k_2 = hf\left(x_n + \frac{h}{2}, y_n + \frac{k_1}{2}\right)$

$$k_3 = hf\left(x_n + \frac{h}{2}, y_n + \frac{k_2}{2}\right); \quad k_4 = hf(x_n + h, y_n + k_3)$$

Step2: Find $y_{n+1} = y_n + \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4)$

Milne's Method:

Step1: Find $f_1 = f(x_1, y_1), f_2 = f(x_2, y_2)$ and $f_3 = f(x_3, y_3).$

Step2: Predictor Formula $y_4^{(P)} = y_0 + \frac{4h}{3}[2f_1 - f_2 + 2f_3]$

Step3: Compute $f_4 = f(x_4, y_4^{(P)})$

Step4: Corrector Formula $y_4^c = y_2 + \frac{h}{3}(f_2 + 4f_1 + f_4^{(P)})$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

Adams – Bashforth Method:

Step1: Find $f_0 = f(x_0, y_0)$, $f_1 = f(x_1, y_1)$, $f_2 = f(x_2, y_2)$ and $f_3 = f(x_3, y_3)$.

Step2: Predictor Formula $y_4^P = y_3 + \frac{h}{24} (55f_3 - 59f_2 + 37f_1 - 9f_0)$

Step3: Compute $f_4 = f(x_4, y_4^P)$

Step4: : Corrector Formula $y_4^c = y_3 + \frac{h}{24} (f_1 - 5f_2 + 19f_1 - 9f_4^P)$

Modular Arithmetics:

Set of Natural Numbers (N): {1, 2, 3,}

Set of Integers (I) = set of natural, negative naturals and zero: $N \cup -N \cup \{0\}$

Greatest common divisor: g or $d = gcd(a, b)$

Relative Primes: g or $d = gcd(a, b) = 1$

Division Algorithm: For integers a and b , with $a > 0$, there exist integers q and r

such that $b = q \cdot a + r$ and $0 \leq r < a$, where q : quotient, r : remainder

6. Euclidean Algorithm

b	a	$b = q_1 a + r_1$	$r_1 = b - a q_1$
a	r_1	$a = q_2 r_1 + r_2$	$r_2 = a - q_2 r_1$

$g = d \cdot gcd(a, b) = ax + by$ with x and y integers

7. Modular and modular class: $m/(a - b)$ then $a \equiv b(modm)$, $0 \leq |b| < m$

$b \equiv a(modm)$, $0 \leq |a| < m$

If $a \equiv b(modm)$ then $a - b = k \cdot m$

8. Properties of modular arithmetic:

- $a \equiv a(modm)$
- $a \equiv b(modm)$ and $b \equiv c(modm)$, then $a \equiv c(modm)$
- $a \equiv c(modm)$ and $b \equiv c(modm)$, then $a \equiv b(modm)$
- $a \equiv b(modm)$ then for any c , $a + c \equiv (b + c)(modm)$





VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

- If $a \equiv b \pmod{m}$ then for any c , $a + c \equiv (b + c) \pmod{m}$
- If $a \equiv b \pmod{m}$ then for any c , $a \cdot c \equiv (b \cdot c) \pmod{m}$
- If $ac \equiv (bc) \pmod{m}$ and $d = \gcd(m, c)$, then $a \equiv b \pmod{m/d}$
- If $ac \equiv (bc) \pmod{m}$ and m is a prime number, then $a \equiv b \pmod{m}$
- If $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then $a \pm c \equiv (b \pm d) \pmod{m}$
- If $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then $a \cdot c \equiv (b \cdot d) \pmod{m}$
- If $ac \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then $ab \equiv b \pmod{m}$
- If $a \equiv b \pmod{m}$, then $a^n \equiv b^n \pmod{m}$
- If $\gcd(m, n) = 1$, $a \equiv b \pmod{m}$ and $a \equiv b \pmod{n}$, then $a \equiv b \pmod{mn}$
- If $ab \equiv 1 \pmod{n}$, then a is inverse of b and b is the inverse of a

Linear Diophantine equation:

$$ax + by = c, \text{ with } d = \gcd(a, b) \text{ has solution if } d \mid c$$

(i) has d congruent solutions and if x_0 and y_0 are primitive solutions,

$$\text{then } x = x_0 + \left(\frac{b}{d}\right)t, \quad y = y_0 + \left(\frac{a}{d}\right)t, \text{ for positive integer } t$$

(ii) if $d = 1$ then it has unique solution

Remainder theorem:

$$x \equiv a_i \pmod{m_i}, \quad i = 1, 2, 3, \text{ with } (m_i, m_j) = 1, \quad i \neq j$$



Procedure to apply remainder theorem

If $m_i, i = 1, 2, 3$ are relatively prime

Then the solution of $x_i \equiv b_i \pmod{m_i}$, $i = 1, 2, 3$ is $x = \sum_1^3 b_i M_i y_i$

where, $M_i = M/m_i$ with $M = \prod_1^3 m_i$

$M_i y_i \equiv 1 \pmod{m_i}$ where y_i is inverse of M_i under mod m_i

11. Fermat's little theorem:

If p is any prime and $p \nmid a$, then $a^{p-1} \equiv 1 \pmod{p}$ or $a^p \equiv a \pmod{p}$

12. Euler's Φ function:

If n is any composite number with prime factorization $n = p_1^{m_1} \times p_2^{m_2} \times \dots$; then number of primes up to n is

$$\Phi(n) = n \left(1 - \frac{1}{p_1}\right) \left(1 - \frac{1}{p_2}\right) \dots \dots$$

13. Wilson's theorem: Prime p divides $(p-1)! + 1$



VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI

MATHEMATICS HANDBOOK

14. RSA cryptosystem:

If p and q are large prime numbers and a is any integer then,
plain text M is encrypted to c by, $c \equiv M^e \pmod{pq}$

The public key will be $= (pq, e)$

d the decryption key the inverse of e , then $d \cdot e \equiv 1 \pmod{(p-1)(q-1)}$
then $M \equiv c^d \pmod{pq}$.

Curvilinear Coordinates:

Curvilinear coordinates are often used to define the location or distribution of physical quantities which may be scalars, vectors or tensors.

If $\vec{R} = x(\mathbf{u}, \mathbf{v}, \mathbf{w})\hat{i} + y(\mathbf{u}, \mathbf{v}, \mathbf{w})\hat{j} + z(\mathbf{u}, \mathbf{v}, \mathbf{w})\hat{k}$, then

- (i) **Tangent vectors** to the u-curve, the v-curve and the w-curve are $\frac{\partial \vec{R}}{\partial u}$, $\frac{\partial \vec{R}}{\partial v}$ and $\frac{\partial \vec{R}}{\partial w}$ respectively
- (ii) **Scale factors** $h_1 = \left| \frac{\partial \vec{R}}{\partial u} \right|$, $h_2 = \left| \frac{\partial \vec{R}}{\partial v} \right|$ and $h_3 = \left| \frac{\partial \vec{R}}{\partial w} \right|$
- (iii) **Unit Tangent Vectors** to the u-curve, the v-curve and the w-curve are

$$T_u = \frac{\left(\frac{\partial \vec{R}}{\partial u} \right)}{h_1}, T_v = \frac{\left(\frac{\partial \vec{R}}{\partial v} \right)}{h_2} \text{ and } T_w = \frac{\left(\frac{\partial \vec{R}}{\partial w} \right)}{h_3}, \text{ respectively.}$$

Normal to the surfaces $\mathbf{u} = \mathbf{u}_0$, $\mathbf{v} = \mathbf{v}_0$ and $\mathbf{w} = \mathbf{w}_0$ are

$$\nabla \mathbf{u} = \frac{T_u}{h_1}, \nabla \mathbf{v} = \frac{T_v}{h_2} \text{ and } \nabla \mathbf{w} = \frac{T_w}{h_3} \text{ respectively}$$

Unit normal vectors to the surfaces $\mathbf{u} = \mathbf{u}_0$, $\mathbf{v} = \mathbf{v}_0$ and $\mathbf{w} = \mathbf{w}_0$ are

$$N_u = \frac{\nabla \mathbf{u}}{|\nabla \mathbf{u}|}, N_v = \frac{\nabla \mathbf{v}}{|\nabla \mathbf{v}|} \text{ and } N_w = \frac{\nabla \mathbf{w}}{|\nabla \mathbf{w}|}, \text{ respectively}$$

Polar Coordinates (r, θ) :

Coordinate transformations: $x = r \cos \theta$, $y = r \sin \theta$

Jacobian: $\frac{\partial(x,y)}{\partial(r,\theta)} = r$

$$(Arc - length)^2: (ds)^2 = (dr)^2 + r^2(d\theta)^2$$






VISVESVARAYA TECHNOLOGICAL UNIVERSITY, BELAGAVI MATHEMATICS HANDBOOK

Cylindrical Coordinates (ρ, φ, z) :

Coordinate transformations: $x = \rho \cos \varphi$, $y = \rho \sin \varphi$, $z = z$




Jacobian: $\frac{\partial(x,y,z)}{\partial(\rho,\varphi,z)} = \rho$

(Arc - length) 2 : $(ds)^2 = (d\rho)^2 + \rho^2(d\varphi)^2 + (dz)^2$

Volume element: $dV = \rho d\rho d\varphi dz$

Spherical Polar Coordinates (r, θ, φ) :

Coordinate transformations: $x = r \sin \theta \cos \varphi$, $y = r \sin \theta \sin \varphi$, $z = r \cos \theta$



Jacobian: $\frac{\partial(x,y,z)}{\partial(r,\theta,\varphi)} = r^2 \sin \theta$

(Arc - length) 2 : $(ds)^2 = (dr)^2 + r^2(d\theta)^2 + (r \sin \theta)^2(d\varphi)^2$

Volume element: $dV = r^2 \sin \theta dr d\theta d\varphi$

