# Third Semester B.E. Degree Examination, June/July 2023 **Engineering Mathematics – III**

Time: 3 hr

Max. Marks: 100

Note: Answer any FIVE full questions, choosing ONE full question from each module.

### Module-1

Expand  $f(x) = x - x^2$  as a Fourier series in the interval  $(-\pi, \pi)$ .

(08 Marks)

- Obtain the half range Fourier cosine series for the function  $f(x) = \sin x$ ,  $0 \le x \le \pi$ . (06 Marks)
- Obtain the constant term and the coefficient of the first sine and cosine terms in the Fourier series of f(x) as given in the following table:

X	0	1	2	3	4	5
f(x)	9	18	24	28	26	20

(06 Marks)

- Obtain the Fourier series for  $f(x) = \sin mx$  in the range  $(-\pi, \pi)$  where m is neither zero nor an (08 Marks)
  - Obtain half range cosine series for

$$f(x) = \begin{cases} Kx, & 0 \le x \le \ell/2 \\ K(\ell - x), & \ell/2 \le x \le \ell \end{cases}$$

 $f(x) = \begin{cases} Kx, & 0 \le x \le \ell/2 \\ K(\ell-x), & \ell/2 \le x \le \ell \end{cases}$  and hence deduce that  $\frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots = \frac{\pi^2}{8}$ 

(06 Marks)

Express Y as a Fourier series upto first harmonic, given that

X	0	$\frac{\pi}{3}$	$\frac{^{\prime }2\pi }{3}$	π	$\frac{4\pi}{3}$	$\frac{5\pi}{3}$	2π
Y	7.9	7.2	3.6	0.5	0.9	6.8	7.9

(06 Marks)

## Module-2

Hence evaluate Fourier transform

$$\int_{0}^{\infty} \frac{x \cos x - \sin x}{x^{3}} \cos \left(\frac{x}{2}\right) dx.$$

(08 Marks)

b. Obtain the Fourier cosine transform of e-ax

(06 Marks)

(06 Marks)

c. Obtain the inverse Z-transform of 
$$\frac{2z^2 + 3z}{(z+2)(z-4)}$$
.

### OR

4 a. Given 
$$Z(u_n) = \frac{2z^2 + 3z + 4}{(z-3)^3}$$
,  $|z| > 3$ , show that  $u_0 = 0$ ,  $u_1 = 2$ ,  $u_2 = 21$ . (08 Marks)

b. Solve by using Z-transforms, 
$$y_{n+2} + 6y_{n+1} + 9y_n = 2^n$$
 with  $y_0 = y_1 = 0$ . (06 Marks)

Find the Fourier transform of  $f(x) = e^{-|x|}$ (06 Marks) Module-3

5 a. Find the coefficient of correlation for the following data:

X	50	50	55	60	65	65	65	60	60	50
у	11	13	14	16	16	15	15	14	13	13

(08 Marks)

b. By the method of least square, find the straight line that best fits the following data:

X	1	2	3	4	5
у	14	27	40	55	68

(06 Marks)

c. Use Newton-Raphson method to find a root of the equation  $\tan x - x = 0$  near x = 4.5 (x is in radians) carry out three iterations. (06 Marks)

OR

6 a. Obtain the lines of regression and hence find the coefficient of correlation for the data:

X	1	2	3	4	5	6	7
У	9	8	10	12	11	13	14

(08 Marks)

b. Fit a second degree parabola to the following data:

X	1	2	3	4	5
у	10	12	13	16	19

(06 Marks)

c. Find the real root of the equation  $xe^x - 3 = 0$  by Regula-Falsi method, correct to three decimal places in (1, 2).

Module-4

7 a. From the following table find f(86) using Newton's backward interpolation formula:

X	40	50	<b>60</b>	70	80	90
f(x)	180	204	226	250	276	304

(08 Marks)

b. Given the values:

X	5	7	11	13	17
f(x)	150	392	1452	2360	5202

Evaluate f(9), using Newton's divided difference formula.

(06 Marks)

c. Compute the value of  $\int_{0.2}^{1.4} (\sin x - \log x + e^x) dx$  using Simpson's  $\frac{3}{8}$  rule taking six equal parts.

OR

8 a. Find an approximate value of f(x) at x = 1.1 from the following data:

X	1	1.2	1.4	1.6	1.8	2
f(x)	0	0.128	0.544	1.296	2.432	4

CMRIT LIBRARY BANGALORE - 560 037

(08 Marks)

b. Find the polynomial f(x) by using Langrage's formula from the following data:

X	0	1	2	5
f(x)	2	3	12	147

(06 Marks)

c. Evaluate  $\int_{0}^{5.2} \log_e x dx$  by Weddle's rule taking six equal strips.

(06 Marks)

Module-5

- 9 a. Verify Green's theorem for  $\oint_C (xy + y^2)dx + x^2dy$  where c is the closed curve of the region bounded by y = x and  $y = x^2$ . (08 Marks)
  - b. Evaluate  $\int_{C}^{C} xy dx + xy^{2} dy$  by Stoke's theorem where c is the square in the x-y plane with vertices (1, 0) (-1, 0) (0, 1) (0, -1).
  - c. Derive Euler's equation  $\frac{\partial f}{\partial y} \frac{d}{dx} \left[ \frac{\partial f}{\partial y'} \right] = 0$  (06 Marks)

OR

CMRIT LIBRARY

- 10 a. If  $\vec{F} = (2x^2 3z)\hat{i} 2xy\hat{j} 4x\hat{k}$ , evaluate  $\iiint_{V} \nabla \cdot \vec{F} dv$  where v is the region bounded by the planes x = 0, y = 0, z = 0 and 2x + 2y + z = 4. (08 Marks)
  - b. Find the extremal of the functional  $\int_{0}^{x_2} [(y')^2 y^2 + 2y \sec x] dx$ . (06 Marks)
  - c. Prove that the shortest distance between two points in a plane is along the straight line joining them. (06 Marks)