



CBCS SCHEME

USN

18MAT31

Third Semester B.E. Degree Examination, June/July 2024 Transform Calculus, Fourier Series and Numerical Techniques

Time: 3 hrs.

Max. Marks: 100

Note: Answer any FIVE full questions, choosing ONE full question from each module.

Module-1

- 1 a. Find the Laplace transform of
 i) $e^{-t} \cos^2 3t$ ii) $t \cos t$

(06 Marks)

- b. A periodic function of period $\frac{2\pi}{\omega}$ is defined by

$$f(t) = \begin{cases} E \sin \omega t, & 0 \leq t \leq \frac{\pi}{\omega} \\ 0, & \frac{\pi}{\omega} \leq t \leq \frac{2\pi}{\omega} \end{cases} \quad \text{where } E \text{ and } \omega \text{ are constants.}$$

$$\text{Show that } L\{f(t)\} = \frac{E\omega}{(s^2 + \omega^2)(1 - e^{-\pi s/\omega})}$$

(07 Marks)

- c. Find the Inverse Laplace transform of

$$\text{i) } \frac{2s-1}{s^2+2s+17} \quad \text{ii) } \log\left(\frac{s^2+1}{s(s+1)}\right)$$

(07 Marks)

OR

- 2 a. Express the function $f(t)$ in terms of unit step function and find its Laplace transform, where

$$f(t) = \begin{cases} \cos t, & 0 < t \leq \pi \\ 1, & \pi < t \leq 2\pi \\ \sin t, & t > 2\pi \end{cases}$$

(06 Marks)

- b. Using the convolution theorem, obtain inverse Laplace transform of $\frac{s}{(s+1)(s^2+1)}$

(07 Marks)

- c. Solve the equation $y'' + 5y' + 6y = e^t$ under the condition $y(0) = 0, y'(0) = 0$

(07 Marks)

Module-2

- 3 a. Find the Fourier series of the function $f(x) = x^2$ in $(-\pi, \pi)$.
 b. Define half range sine and cosine series in the interval $(0, l)$.
 c. Find the constant term and the first two harmonics in the fourier series for $f(x)$ given by the following table.

(08 Marks)

(04 Marks)

x	0	$\pi/3$	$2\pi/3$	π	$4\pi/3$	$5\pi/3$	2π
$f(x)$	1.0	1.4	1.9	1.7	1.5	1.2	1.0

(08 Marks)

OR

- 4 a. Obtain the fourier series of the saw-tooth function

$$f(x) = \frac{Ex}{T} \quad \text{for } 0 < x < T \quad \text{given that } f(x+T) = f(x) \quad \text{for all } x > 0. \quad (06 \text{ Marks})$$

- b. Obtain the Fourier series expansion of

$$f(x) = \begin{cases} \pi x & \text{in } 0 \leq x \leq 1 \\ \pi(2-x) & \text{in } 1 \leq x \leq 2 \end{cases} \quad \text{over the interval } (0, 2)$$

$$\text{Deduce that } \frac{\pi^2}{8} = \frac{1}{1^2} + \frac{1}{3^2} + \frac{1}{5^2} + \dots \quad (07 \text{ Marks})$$

- c. Expand $f(x) = \sin x$ in half range cosine series over the interval $(0, \pi)$. (07 Marks)

Module-3

- 5 a. Prove that fourier transform of

$$f(x) = \begin{cases} 1 + \frac{x}{a}, & -a < x < 0 \\ 1 - \frac{x}{a}, & 0 < x < a \\ 0, & \text{otherwise} \end{cases} \quad \text{is } \frac{4 \sin^2 \frac{au}{2}}{au^2}, \quad \text{if Fourier transform of } f(x) \text{ is } F(u). \quad (06 \text{ Marks})$$

- b. Find the Fourier sine transform of $f(x) = e^{-|x|}$ and hence

$$\text{evaluate } \int_0^\infty \frac{x \sin mx}{1+x^2} dx, \quad m > 0. \quad (07 \text{ Marks})$$

$$\text{c. Find z-transform of } 5n^2 + 4 \sin\left(\frac{n\pi}{2} + \frac{\pi}{4}\right) \quad (07 \text{ Marks})$$

OR

- 6 a. Find the fourier cosine transform of

$$f(x) = \begin{cases} x & \text{for } 0 < x < 1 \\ 2-x & \text{for } 1 < x < 2 \\ 0 & \text{for } x > 2 \end{cases} \quad \text{CMRIT LIBRARY BANGALORE - 560 037} \quad (07 \text{ Marks})$$

$$\text{b. Obtain the inverse z-transform of } \frac{4z^2 - 2z}{(z-1)(z-2)^2} \quad (07 \text{ Marks})$$

- c. Solve the difference equation

$$u_{n+2} + 3u_{n+1} + 2u_n = 3^n, \quad \text{given } u_0 = 0, \quad u_1 = 1, \quad \text{using z-transform.} \quad (06 \text{ Marks})$$

Module-4

- 7 a. Use Taylor's series method to find the value of y at $x = 0.1$, given that $dy/dx = x^2 + y^2$, $y(0) = 1$. Consider upto 4th degree term. (06 Marks)

- b. By using modified Euler's method, solve the initial value problem $\frac{dy}{dx} = \log(x+y)$, $y(1) = 2$ at the point $x = 1.2$. Take $h = 0.2$ and carryout two modifications. (07 Marks)

- c. Given $\frac{dy}{dx} = xy + y^2$, $y(0) = 1$, $y(0.1) = 1.1169$, $y(0.2) = 1.2773$, $y(0.3) = 1.5049$. Find $y(0.4)$ correct to three decimal places using Milne's predictor – corrector method. Apply corrector formula once. (07 Marks)

OR

- 8 a. Using modified Euler's method compute $y(1.1)$ correct to five decimal places taking $h = 0.1$, given that $\frac{dy}{dx} + \frac{y}{x} = \frac{1}{x^2}$ and $y = 1$ at $x = 1$. (06 Marks)
- b. Use fourth order Runge-Kutta method to find y at $x = 0.1$, given that $\frac{dy}{dx} = 3e^x + 2y$, $y(0) = 0$ and $h = 0.1$. (07 Marks)
- c. Apply Adam's - Bashforth method to solve the equation $(y^2 + 1)dy - x^2 dx = 0$ at $x = 1$ given $y(0) = 1$, $y(0.25) = 1.0026$, $y(0.5) = 1.0206$, $y(0.75) = 1.0679$. Apply corrector formula once. (07 Marks)

Module-5

- 9 a. By Runge-Kutta method solve $y'' = xy'^2 - y^2$ for $x = 0.2$ correct to four decimal places, using initial conditions $y = 1$ and $y' = 0$ when $x = 0$. Take step length $h = 0.2$. (06 Marks)
- b. Derive the Euler's equation in the form $\frac{\partial f}{\partial y} - \frac{d}{dx} \left(\frac{\partial f}{\partial y'} \right) = 0$. (07 Marks)
- c. Prove that geodesics on a plane are straight line. (07 Marks)

CMRIT LIBRARY
BANGALORE - 560 037

OR

- 10 a. Using Runge-Kutta method solve the differential equation at $x = 0.1$ under the given conditions:
- $$\frac{d^2y}{dx^2} = x^3 \left(y + \frac{dy}{dx} \right), \quad y(0) = 1, \quad y'(0) = 0.5. \quad \text{Take step length } h = 0.1. \quad (06 \text{ Marks})$$
- b. Apply Milne's method to compute $y(0.8)$ given that $\frac{d^2y}{dx^2} = 1 - 2y \frac{dy}{dx}$ and the following table of initial values.

x	0	0.2	0.4	0.6
y	0	0.02	0.0795	0.1762
y'	0	0.1996	0.3937	0.5689

Apply corrector formula once. (07 Marks)

- c. Find the extremal of the functional $\int_a^b (x^2 y'^2 + 2y^2 + 2xy) dx$ (07 Marks)
