


CBCS SCHEME



--	--	--	--	--	--	--	--

18MATDIP31

Third Semester B.E. Degree Examination, June/July 2024 Additional Mathematics – I

Time: 3 hrs.

Max. Marks: 100

Note: Answer any FIVE full questions, choosing ONE full question from each module.

Module-1

- 1 a. Show that $(1 + \cos\theta + i\sin\theta)^n + (1 + \cos\theta - i\sin\theta)^n = 2^{n+1} \cos^n\left(\frac{\theta}{2}\right) \cdot \cos\left(\frac{n\theta}{2}\right)$ (07 Marks)
- b. Express $\sqrt{3} + i$ in the polar form and hence find its modulus and amplitude. (07 Marks)
- c. Find the argument of $\frac{1+i\sqrt{3}}{1-i\sqrt{3}}$ (06 Marks)

OR

- 2 a. If $\vec{A} = i + 2j + 3k$, $\vec{B} = -i + 2j + k$ and $\vec{C} = 3i + j$, find P such that $\vec{A} + P\vec{B}$ is perpendicular to \vec{C} . (07 Marks)
- b. Find the area of the parallelogram whose adjacent sides are the vectors $\vec{A} = 2i + 4j - 5k$ and $\vec{B} = i + 2j + 3k$. (06 Marks)
- c. If $\vec{A} = 4i + 3j + k$ and $\vec{B} = 2i - j + 2k$, find a unit vector N form a right handed system. (07 Marks)

Module-2

- 3 a. Obtain the Maclaurin's series expansion of $\sin x$ up to term containing x^4 . (07 Marks)
- b. If $U = \sin^{-1}\left[\frac{x^2 + y^2}{x - y}\right]$, prove that $x \frac{\partial U}{\partial x} + y \frac{\partial U}{\partial y} = \tan U$. (07 Marks)
- c. If $U = f(x - y, y - z, z - x)$ prove that $\frac{\partial U}{\partial x} + \frac{\partial U}{\partial y} + \frac{\partial U}{\partial z} = 0$. (06 Marks)

OR

- 4 a. Prove that $\log(1 + x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$ by using Maclaurin's series notation. (07 Marks)
- b. Using Euler's theorem prove that $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 4u \log u$, if $u = e^{\left(\frac{x^3 y^3}{x^2 + y^2}\right)}$ (07 Marks)
- c. If $u = x + y$, $v = y + z$ and $w = z + x$ then find $J\left(\frac{u, v, w}{x, y, z}\right)$. (06 Marks)

Important Note : 1. On completing your answers, compulsorily draw diagonal cross lines on the remaining blank pages.
2. Any revealing of identification, appeal to evaluator and /or equations written eg, 42+8 = 50, will be treated as malpractice.

Module-3

- 5 a. A particle moves along a curve $x = e^{-t}$, $y = 2\cos 3t$ and $z = 2\sin 3t$, where t is the time variable. Determine the components of velocity and acceleration vectors at $t = 0$ in the direction of $i + j + k$. (07 Marks)
- b. Find the unit normal to the surface $x^2y + 2xz = 4$ at $(2, -2, 3)$. (06 Marks)
- c. Show that the vector field $\vec{F} = (4xy - z^3)i + (2x^2)j - (3xz^2)k$ is irrotational. (07 Marks)

OR

- 6 a. Find $\text{div } \vec{F}$ and $\text{curl } \vec{F}$ where $\vec{F} = \nabla(x^3 + y^3 + z^3 - 3xyz)$. (07 Marks)
- b. If $\vec{F} = (3x^2y - z)i + (xz^3 + y^4)j - 2x^3z^2k$, find $\text{grad}(\text{div } \vec{F})$ at $(2, -1, 0)$. (07 Marks)
- c. Find the value 'a' such that the vector field $\vec{F} = (x + 3y)i + (y - 2z)j + (x + az)k$ is Solenoidal. (06 Marks)

Module-4

- 7 a. Obtain the reduction formula for $\int_0^{\pi/2} \cos^n x \, dx$, $n > 0$. (07 Marks)
- b. Evaluate $\int_0^1 \frac{x^9}{\sqrt{1-x^2}} \, dx$ (06 Marks)
- c. Evaluate $\iint_C xy(x+y) \, dx \, dy$ over the area between $y = x^2$ and $y = x$. (07 Marks)

OR

- 8 a. Obtain the reduction formula for $\int_0^{\pi/2} \sin^n x \, dx$, $n > 0$ (07 Marks)
- b. Evaluate $\int_0^{\infty} \frac{x^2}{(1+x^6)^{7/2}} \, dx$ (06 Marks)
- c. Evaluate $\int_0^1 \int_0^{\sqrt{1-x^2}} \int_0^{\sqrt{1-x^2-y^2}} \frac{dx \, dy \, dz}{\sqrt{1-x^2-y^2-z^2}}$ (07 Marks)

Module-5

- 9 a. Solve $(4xy + 3y^2 - x) \, dx + x(x + 2y) \, dy = 0$ (07 Marks)
- b. Solve $\frac{dy}{dx} + \frac{y}{x} = y^2x$ (06 Marks)
- c. Obtain the solution of the differential equation

$$(1 + e^{x/y}) \, dx + e^{x/y} \left(1 - \frac{x}{y}\right) \, dy = 0$$

CMRIT LIBRARY
OR BANGALORE - 560 037

- 10 a. Solve : $\tan y \, dy = (\cos y \cos^2 x - \tan x) \, dx$ (07 Marks)
- b. Solve : $\left[y \left(1 + \frac{1}{x}\right) + \cos y \right] \, dx + (x + \log x - x \sin y) \, dy = 0$ (07 Marks)
- c. Solve : $(1 + y^2) \, dx = (\tan^{-1} y - x) \, dy$ (06 Marks)
