



CBCS SCHEME

21MAT31

Third Semester B.E. Degree Examination, June/July 2024

Transform Calculus, Fourier Series & Numerical Techniques

Time: 3 hrs.

Max. Marks: 100

Note: Answer any FIVE full questions, choosing ONE full question from each module.

Module-1

- 1 a. Find the Laplace Transform of, $\left(\frac{4t+5}{e^{2t}} \right)^2$. (06 Marks)
- b. The square wave function $f(t)$ with period $2a$ is defined by,

$$f(t) = t ; 0 \leq t \leq a$$

$$= 2a - t ; a \leq t \leq 2a$$
Find $L[f(t)]$. (07 Marks)
- c. Evaluate $L^{-1} \left[\frac{s^2}{(s^2 + a^2)^2} \right]$ by applying convolution theorem. (07 Marks)

OR

- 2 a. Find inverse Laplace transform $\frac{2s^2 - 6s + 5}{s^3 - 6s^2 + 11s - 6}$. (06 Marks)
- b. Express the following function in terms of unit step function and hence find the Laplace transform.

$$f(t) = 1 ; 0 < t \leq 1$$

$$= t ; 1 \leq t \leq 2$$

$$= t^2 ; t > 2$$
 (07 Marks)
- c. Applying Laplace transform, solve the differential equation,
 $y''(t) + 4y'(t) + 4y(t) = e^{-t}$,
Subject to the condition $y(0) = y'(0) = 0$. (07 Marks)

Module-2

- 3 a. Obtain the Fourier series of $f(x) = x^2$ over the interval $[-\pi, \pi]$, hence deduce that

$$\frac{\pi^2}{12} = \frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} - \frac{1}{4^2} + \dots + \infty$$
. (06 Marks)
- b. Obtain the half range sine series of the function, $f(x) = x$ in the interval $(0, 2)$. (07 Marks)
- c. Obtain the constant term and co-efficient of first cosine and sine terms in the expansion of y from the following table :

x	0°	60°	120°	180°	240°	300°	360°
y	7.9	7.2	3.6	0.5	0.9	6.8	7.9

(07 Marks)

OR

- 4 a. Find the Fourier series of $f(x) = 2 - x ; 0 \leq x \leq 4$

$$x - 6 ; 4 \leq x \leq 8$$
 (06 Marks)
- b. Obtain the half range sine series of the function, $f(x) = x^2$ over $(0, \pi)$. (07 Marks)

- c. Obtain a_0, a_1, b_1 in the Fourier expansion of y using harmonic analysis for the data given,

x	0	1	2	3	4	5
y	9	18	24	28	26	20

(07 Marks)

Module-3

- 5 a. Find the Fourier sine and cosine transforms of $f(x) = e^{-\alpha x}; \alpha > 0.$ (06 Marks)
- b. Obtain the inverse z-transform of, $\frac{2z^2 + 3z}{(z^2 - 2z - 8)}.$ (07 Marks)
- c. Find the Fourier transform of,

$$f(x) = x^2; |x| < a$$

$$= 0; |x| > a$$
where a is +ve constant. (07 Marks)

OR

- 6 a. Find the Complex Fourier transform of the function,

$$f(x) = 1 \text{ for } |x| \leq a$$

$$= 0 \text{ for } |x| > a$$
Hence deduce, evaluate $\int_0^\infty \frac{\sin x}{x} dx.$ (06 Marks)
- b. Evaluate $Z_T \left[2n + \sin \left(\frac{n\pi}{4} \right) + 1 \right].$ (07 Marks)
- c. Solve the difference equation, $y_{n+2} + 6y_{n+1} + 9y_n = 2^n$ with $y_0 = y_1 = 0$ using Z-Transform. (07 Marks)

Module-4

- 7 a. Classify the following partial differential equation,
- (i) $\frac{\partial^2 u}{\partial x^2} + 4 \frac{\partial^2 u}{\partial x \partial y} + 4 \frac{\partial^2 u}{\partial y^2} - \frac{\partial u}{\partial x} + 2 \frac{\partial u}{\partial y} = 0.$
- (ii) $x^2 \frac{\partial^2 u}{\partial x^2} + (1-y^2) \frac{\partial^2 u}{\partial y^2} = 0, -\infty < x < \infty, -1 < y < 1.$
- (iii) $(1+x^2) \frac{\partial^2 u}{\partial x^2} + (5+2x^2) \frac{\partial^2 u}{\partial x \partial t} + (4+x^2) \frac{\partial^2 u}{\partial t^2} = 0$
- (iv) $(x+1) \frac{\partial^2 u}{\partial x^2} - 2(x+2) \frac{\partial^2 u}{\partial x \partial y} + (x+3) \frac{\partial^2 u}{\partial y^2} = 0$ **CMRIT LIBRARY**
BANGALORE - 560 037 (10 Marks)
- b. Find the numerical solution of the parabolic equation $\frac{\partial^2 u}{\partial x^2} = 2 \frac{\partial u}{\partial t}$, using Schmidt formula.
Given $u(0, t) = 0 = u(4, t)$ and $u(x, 0) = x(4-x)$ by taking $h = 1$ find the values upto $t = 5.$ (10 Marks)

OR

- 8 a. Solve $u_{xx} + u_{yy} = 0$ in the following square region with the boundary conditions as indicated in the Fig. Q8 (a). (10 Marks)




Fig. Q8 (a)

- b. Solve numerically $u_{xx} = 0.0625 u_{tt}$, subject to the conditions $u(0, t) = 0 = u(5, t)$, $u(x, 0) = x^2(x - 5)$ and $u_t(x, 0) = 0$ by taking $h = 1$ for $0 \leq t \leq 1$. (10 Marks)

Module-5

- 9 a. Use Runge-Kutta method to find $y(0.2)$ for the equation, $\frac{d^2y}{dx^2} - x \frac{dy}{dx} - y = 0$. Given that $y=1$, $y'=0$ when $x=0$. (06 Marks)
- b. Find the curves on which the function, $\int_0^1 \{(y')^2 + 12xy\} dx$ with $y(0) = 0$ and $y(1) = 1$ can be extremised. (07 Marks)
- c. Derive the Eulers equation in the form $\frac{\partial f}{\partial y} - \frac{d}{dx} \left(\frac{\partial f}{\partial y'} \right) = 0$ (07 Marks)
- OR**
- 10 a. Solve the differential equation $y'' + xy' + y = 0$ for $x = 0.4$, using Milne's predictor-corrector formula given that,

x	0	0.1	0.2	0.3
y	1	0.995	0.9802	0.956
$\frac{dy}{dx}$	0	-0.0995	-0.196	-0.2863

CMRIT LIBRARY
BANGALORE - 560 037

- b. Find the curve on which functional $\int_0^{\frac{\pi}{2}} [(y')^2 - y^2 + 2xy] dx$ with $y(0) = y\left(\frac{\pi}{2}\right) = 0$ can be extremized. (07 Marks)
- c. Prove that shortest distance between two points in a plane is a straight line. (07 Marks)
